

Hypocrea subpurpurea sp. nov. (Fig. 5)

Stromata extensa, carnosa, subpurpurea, himmifina in sicco, 1.5-2.5 cm² in area, 0.6-0.9 mm crassa, peritheciis fere nudis ex stromatibus, sed efferentibus hyphas pilosis, ellipsoideis, 160-200 μ in diametro verticali, 80-120 μ latis, ostiolis protuberans, collis 20-40 μ longis. Partosporae hyalinae, verrucosae, partosporis distalibus subgloboesis vel obovatis, 2.5-3.5 \times 2.2-2.8 μ ; partosporis proximalibus obovato-ellipsoideis, 3-3.8 \times 2-2.5 μ . St. conid. incognitis.

Habitat: In foris emortuis vel ramis.

Str. effuse, fleshy, pale purple in fresh material, becoming fawn-colored in dried specimens, up to 1.5 \times 2.5 cm², 0.6-0.9 mm thick. Inn. t. of str. close to *t. intricata*, but almost lacking intercellular spaces; cells thin-walled, 4-10 μ in diam.; interperithecial surface filled with parallel hairy hyphae; hyphae 2-3 μ in diam. Perith. almost naked on str., with short hairy hyphae, elliptical with a projecting neck, 160-200 \times 80-120 μ ; necks of ost. 20-40 μ long. P. sp. hyaline, roughly warted; the distal subglobose to obovate, 2.5-3.5 \times 2.2-2.8 μ ; the proximal obovate ellipsoid, 3-3.8 \times 2-2.5 μ . Culture difficult even with fresh and well-matured p.sp.; st.conid. unknown.

Hab. On dead leaves or branchlets of *Fagus crenata*.

Holotype: Daimyojin-Zawa, Sugadaira, Nagano Pref., 3-VIII-1965, Y.D., (TNS.D-38=TNS-F-223218)




Fig. 5. *Hypocrea subpurpurea* sp. nov. (TNS.D-38) and *Hypocrea obliterata* sp. nov. (TNS.D-58). A-D, E-H; A,B,E,F, Vertical section of stroma. C,D,G, Ascii and partospores. H, Inner tissue of stroma.

Basionym: *Hypocrea subgen. Homalocrea* SACC., Syll. Fung. 2: 520-536 (1883).

Lectotype species: *Hypocrea citrina* (PERS. ex FR.) FR. (designated here).